
Identification and establishment of type IV interferons and the characterization of interferon-u including its class II cytokine receptors IFN-uR1 and IL-10R2


Supplementary Figures and Tables

Supplementary Figures 1-29 Supplementary Tables 1-6


Supplementary Figure 1




Supplementary Fig. 1. Induction of antiviral ISGs in IFN- υ -overexpressed zebrafish. Embryos (n = 150) were injected at one-cell stage with IFN- υ or empty vector plasmid, and after 24 hours, the mRNA level of ISGs, including gig2 (a), irf1 (b), mxa (c) and viperin (d), was detected by quantitative RT-PCR. The expression of the selected genes was normalized against gapdh and fold changes were calculated relative to control group (empty vector). Data represent mean \pm SEM from three independent experiments. The two-tailed Student's t-test was used to determine the statistical significance, with \star indicating P < 0.05, and $\star\star$ P < 0.01.




Supplementary Fig. 2. Generation of *ifnu* deficiency zebrafish. (a) The target site and sequence information in *ifnu* mutants. Cas9/gRNA system specific to *ifnu* targets on Exon 4 and leads to 17 bp deletion (5-CAGATCCCAAGACGTTC-3), which results in frameshift mutation and premature translation termination to generate a truncated protein with 88 aa. (b) *ifnu* expression in deficiency zebrafish. Zebrafish larvae (5 dpf, n = 33) were infected with GCRV for 24 hours and were collected to extract RNA to determine the expression of *ifnu*, which was normalized against *gapdh* by quantitative RT-PCR. The word, MOCK, represents the zebrafish larvae (6 dpf, n = 33) which were not infected with the virus. Data represent mean \pm SEM from three independent experiments. The two-tailed Student's *t*-test was used to determine the statistical significance, * indicating P < 0.05, and ** P < 0.01.




Supplementary Fig. 3. Effects on viral titre in response to GCRV infection by knockdown of IFN- ϕ 1, IFN- γ and IFN- γ 1 in *ifnu* deficiency zebrafish. Embryos at one-cell stage (n = 150) were micro-injected with the IFN morpholinos or standard (std) control morpholino for 72 hours, and hatched zebrafish larvae (n = 33) were infected with GCRV for 24 hours and then collected to detect viral titers. Data represent mean \pm SEM from three independent experiments. The two-tailed Student's *t*-test was used to determine the statistical significance, with * indicating P < 0.05, and ** P < 0.01.




Supplementary Fig. 4. Knockdown test of CRFBs in zebrafish. Inhibitory effects of zebrafish IFN possible receptors' morpholinos (MOs) on *viperin* (a) or *irgf1* (b) expression induced by IFN1 (IFN- φ 1) / IFN3 (IFN- φ 3) or IFN- φ 1 (IFN- φ 2) / IFN- φ 1), respectively. IFN plasmids and different MOs were co-injected into one-cell stage embryos (n = 150), and at 2 dpf the mRNA level of *viperin* or *irgf1* was detected by quantitative RT-PCR. The expression of selected genes was normalized against *gapdh*, and fold changes were calculated relative to control group which were co-injected into embryos with empty plasmid and standard control MOs. Data represent mean ± SEM from three independent experiments. The two-tailed Student's *t*-test was used to determine the statistical significance, with * indicating *P* < 0.05 and ** indicating *P* < 0.01. Knockdown effect and strategy for the *crfb4*-MO (c), *crfb7*-MO (d), *crfb8*-MO (e), *crfb12*-MO (f), *crfb14*-MO (g), *crfb15*-MO (h) and *crfb16*-MO (i). After the injection of *crfb*-MOs for 48 hours, RT-PCRs and sequencing of PCR production were performed to detect knockdown effect of different MOs. Arrows and solid lines represent PCR primers (Supplementary Table 3) and MOs, respectively. Data represent two independent experiments. All MOs described above have been proved previously to possess knockdown activity¹⁻³.



Supplementary Fig. 5. Effects on IFN- υ -induced ISG expression by knockdown of CRFB morpholinos in zebrafish at 24 hpf. IFN- υ expressing plasmid and different morpholinos were co-injected into zebrafish embryos (n = 500) at one-cell stage, and 24 hours later the expression of ISGs was detected by quantitative RT-PCR. The expression of the selected genes was normalized against *gapdh* and fold changes were calculated relative to control group which was co-injected in embryos with empty plasmid and standard (std.) control morpholinos. Data represent mean \pm SEM from three independent experiments.




Supplementary Fig. 6. Expressional pattern of *ifnur1* in zebrafish embryos. (a) Detection of *ifnur1* expression by whole-mount *in situ* hybridization (WISH). Data represent two independent experiments. (b) Expression of *ifnur1* was detected by quantitative RT-PCR. The expression level of *ifnur1* was calculated relative to *gapdh*. Data represent mean ± SEM from three independent experiments.



Supplementary Fig. 7. Generation of *ifnur1* and *crfb4* deficiency zebrafish. (a, b) Cas9/gRNA system specific to *ifnur1* (a) and *crfb4* (b) targets on Exon 3, and leads to 56 bp (5-


AGGTCCTGACTGTAGGATTATTGGATTTTAAGGCTACAGCTGAATGGCTTCCAGGA-3) and 5 bp (5-CAGTG-3) deletion, respectively, which results in frameshift mutation and premature translation termination. (c, d) Detection of protein expression for CRFB4 (c) and IFNUR1 (d) in deficiency and WT zebrafish by Western blotting. GAPDH protein was detected as control. Data represent two independent experiments.



Supplementary Fig. 8. Observation of embryonic development phenotype for *ifnu*, *ifnur1* (*crfb12*) and *crfb4* knockout at 24 (a), 48 (b) and 72 (c) hpf.


a ifnu^{-/-} crfb12^{-/-} crfb4^{-/-} wt crfb12^{-/-} ifnu^{-/-} crfb4^{-/-} b wt crfb12^{-/-} ifnu^{-/-} crfb4^{-/-} C wt


Supplementary Fig. 9. Observation of embryonic development phenotype for *ifnu*, *ifnur1* (*crfb12*) and *crfb4* knockout at 30 (a), 60 (b) and 90 (c) dpf.




Supplementary Fig. 10. Observation of embryonic development phenotypes for the knockdown of CRFBs and control at 24 hpf. Different morpholinos for these CRFBs (1, 2, 4, 5, 6, 7, 8, 12, 13, 14, 15, 16 and 17) were injected separately into one-cell stage embryos, and 24 hours later embryos were collected for observation, respectively.

Supplementary Figure 11 crfb1 crfb2 crfb4 crfb5 crfb6 crfb7 crfb8 crfb12 crfb13 crfb14 crfb15 crfb16







Supplementary Fig. 11. Observation of embryonic development phenotypes for the knockdown of CRFBs and control 48 hpf. Different morpholinos for these CRFBs (1, 2, 4, 5, 6, 7, 8, 12, 13, 14, 15, 16 and 17) were injected separately into one-cell stage embryos, and 48 hours later embryos were collected for observations, respectively.




Supplementary Fig. 12. Sequence alignment of IFN- υ proteins in vertebrates. The conserved CXXXXX[W/L] sequences are marked in arrows. Identical and similar amino acids are indicated by shade with black and grey, respectively. The intron positions and their phases were indicated by red line.




Supplementary Fig. 13. Sequence alignment (a) and phylogenetic analysis (b) of IFN genes in zebrafish. Identical and similar amino acids are indicated by shade with black and grey, respectively. The putative signal peptides are highlighted by solid lines above the alignment. Neighbor-joining (NJ) method was used to construct the phylogenetic tree.




Supplementary Fig. 14. Sequence alignment (a) and phylogenetic analysis (b) of the IFN genes in clawed frog. Identical and similar amino acids are indicated by shade with black and grey, respectively. The putative signal peptides are highlighted by solid lines above the alignment. Neighbor-joining (NJ) method was used to construct the phylogenetic tree.




Supplementary Fig. 15. Sequence alignment (a) and phylogenetic analysis (b) of IFN genes in green anole. Identical and similar amino acids are indicated by shade with black and grey, respectively. The putative signal peptides are highlighted by solid lines above the alignment. Neighbor-joining (NJ) method was used to construct the phylogenetic tree.




Supplementary Fig. 16. Sequence alignment (a) and phylogenetic analysis (b) of IFN genes in chicken. Identical and similar amino acids are indicated by shade with black and grey, respectively. The putative signal peptides are highlighted by solid lines above the alignment. Neighbor-joining (NJ) method was used to construct the phylogenetic tree.




Supplementary Fig. 17. Sequence alignment (a) and phylogenetic analysis (b) of IFN genes in platypus. Identical and similar amino acids are indicated by shade with black and grey, respectively. The putative signal peptides are highlighted by solid lines above the alignment. Neighbor-joining (NJ) method was used to construct the phylogenetic tree.



Supplementary Fig. 18. Structural comparison of IFNs in zebrafish. (a) The structures of type I (IFN- ϕ 1/IFN1, 3piv and IFN- ϕ 2/IFN2, 3piw) and putative structures of type II (IFNG) and IFNU. Comparison of IFNU to IFN- ϕ 1 (b), IFNG (c) and IFN- ϕ 2 (d). Putative helices of IFNU were indicated by A to E. The overlaid regions on structure were highlighted by orange box in the alignments.




Supplementary Fig. 19. Structural comparison of IFNs in clawed frog. (a) The putative structures of type I (IFN1), type II (IFNG), type III (IFNL) and IFNU. Comparison of IFNU to IFN1 (b), IFNG (c) and IFNL (d). Putative helices of IFNU were indicated by A to E. The overlaid regions on structure were highlighted by orange box in the alignments.




Supplementary Fig. 20. Structural comparison of IFNs in green anole. (a) The putative structures of type I (IFN1), type II (IFNG), type III (IFNL) and IFNU. Comparison of IFNU to IFN1 (b), IFNG (c) and IFNL (d). Putative helices of IFNU were indicated by A to E. The overlaid regions on structure were highlighted by orange box in the alignments.

Supplementary Figure 21 а GG IFNU GG IFNB GG IFNG GG IFNL b С d


Supplementary Fig. 21. Structural comparison of IFNs in chicken. (a) The putative structures of type I (IFNB), type II (IFNG), type III (IFNL) and IFNU. Comparison of IFNU to IFNB (b), IFNG (c) and IFNL (d). Putative helices of IFNU were indicated by A to E. The overlaid regions on structure were highlighted by orange box in the alignments.




Supplementary Fig. 22. Structural comparison of IFNs in platypus. (a) The putative structures of type I (IFN1), type II (IFNG), type III (IFNL3L) and IFNU. Comparison of IFNU to IFN1 (b), IFNG (c) and IFNL3L (d). Putative helices of IFNU were indicated by A to E. The overlaid regions on structure were highlighted by orange box in the alignments.




Supplementary Figure 23 - continued




Supplementary Fig. 23. Sequence alignment of IFN receptor proteins in vertebrates. Identical and similar amino acids are indicated by shade with black and grey, respectively. The putative signal peptide, D200 domain and transmembrane region are highlighted by solid lines above the alignment. The intron positions and phases of receptor genes were indicated by red (IFNUR1), blue (IFNAR2), yellow (IFNLR1), green (IFNAR1) and pink (IL10R2) lines, respectively.




Supplementary Fig. 24. Sequence alignment of extracellular regions in IFN receptor proteins in vertebrates. Identical and similar amino acids are indicated by shade with black and grey, respectively. The putative signal peptide and D200 domain are highlighted by solid lines above the alignment. The helices and beta-sheet related structures of IFNLR1 are indicated by underline with "h" and "b", respectively.




Supplementary Figure 25 - continued




Supplementary Fig. 25. Sequence alignment of intracellular regions in IFN receptor proteins in vertebrates. Identical and similar amino acids are indicated by shade with black and grey, respectively. The conserved peptide motif (PXXL) and a hydrophobic residues-rich sequence of IFNUR1 are indicated by 'box1' and 'box2' above the alignment, respectively. The conserved tyrosine of IFNUR1 is marked by arrows. The known docking sites for JAK1, JAK2, TYK2, STAT1, STAT2 and STAT3 are indicated by dark blue, light blue, green, orange, red and yellow box, respectively.




Supplementary Fig. 26. Collinearity analysis of *ADARB2-PFKP* locus (a) and *CLIC2-BRCC3* locus (b) in mammals. All genes are indicated with arrow symbols which point to the transcription direction. *IFNU* and *IFNUR1* are dyed in red. Gene name abbreviation: WDR37, WD repeat domain 37; ADARB2, adenosine deaminase RNA specific B2 (inactive); PFKP, phosphofructokinase, platelet; PITRM1, pitrilysin metallopeptidase 1; FAAH2, fatty acid amide hydrolase 2; TMLHE, trimethyllysine hydroxylase, epsilon; VAMP7, vesicle associated membrane protein 7; VBP1, VHL binding protein 1; SPRY3, sprouty RTK signaling antagonist 3; RAB39B, RAB39B, member RAS oncogene family; BRCC3, BRCA1/BRCA2-containing complex subunit 3; PICALM, phosphatidylinositol binding clathrin assembly protein; ZXDB, zinc finger X-linked duplicated B.



Supplementary Fig. 27. Gene synteny of *IFNUR1* (a) and other IFN receptor gene (b-d) loci in vertebrates. All genes are indicated with arrow symbols which point to the transcription direction. *IFNUR1* are dyed in red, and the conserved and non-conserved neighbor genes are illustrated in grey and white, respectively. Gene name abbreviation: AIFM1, apoptosis inducing factor mitochondria associated 1; DNAJC28, DnaJ heat shock protein family (Hsp40) member C28; TMEM50B, transmembrane protein 50B; MYOM3, myomesin 3; GRHL3, grainyhead like transcription factor 3; STPG1, sperm tail PG-rich repeat containing 1; NIPAL3, NIPA like domain containing 3; RCAN3, RCAN family member 3; OLIG3, oligodendrocyte transcription factor 3; SLC35D3, solute carrier family 35 member D3.



Supplementary Fig. 28. Phylogenetic analysis of IFNUR1 in vertebrates. Protein sequences from class II cytokine receptors in vertebrates were used to construct the neighbor-joining (NJ) tree.



Supplementary Fig. 29. Antiviral function of *X. laevis* (xl, clawed frog) IFN- υ . (a) Western blotting detection of recombinant flag-tag-IFN- υ protein. Vector and UT represent supernatant media from empty plasmid-transfected and untreated HEK293T cells (negative control), respectively. Data represent two independent experiments. (b) Antiviral ISGs induced by xl-IFN- υ . A6 cells (1.0×10^6) were incubated with control medium and recombinant xl-IFN- υ containing supernatant produced in HEK293T cells for 10 hours. The mRNA expression level of ISGs (mx1 and *viperin*) was determined by quantitative RT-PCR. The expression of the selected genes was normalized against *actb* and fold changes were calculated relative to control group. (c) Viral yields of FV3 reduced by xl-IFN- υ . Recombinant xl-IFN- υ and control medium were added to A6 cells (1.0×10^6) for 10 hours incubation. Subsequently, the cells were infected with FV3 at the multiplicity of infection (MOI) of 5, and viral titre was measured by plaque assay. The knockdown effects of *IFNUR1* (d) and *IL10RB* (e) siRNAs were detected by quantitative RT-PCR. (f-h) Effects on xl-IFN- υ -induced ISG expression and antiviral function by knockdown of *IFNUR1* and *IL10RB*. ISG expression and viral titre were determined by quantitative RT-PCR and plaque assay, respectively, after incubation with xl-IFN- υ and knockdown of *IFNUR1* or *IL10RB* in A6 cells. Data represent mean \pm SEM from three independent experiments. The two-tailed Student's *t*-test was used to determine the statistical significance, \star indicating P < 0.05, and $\star\star P < 0.01$.

Supplementary Table 1.Comparison of class II cytokine receptors between zebrafish and human

Gene in human	Gene in zebrafish	Synonym in fish	Reference
IFNAR1	CRFB5	IFNAR1, IFNphiR2	2, 4
IFNAR2	CRFB1 (duplication)	IFNAR2-1, IFNphi1R1	2, 4
	CRFB2 (duplication)	IFNAR2-2, IFNphi2R1	2, 4
	CRFB3 (duplication, lost)	_	3
IFNGR1	CRFB13 (duplication)	IFNGR1-2	1, 4
	CRFB17 (duplication)	IFNGR1-1	1, 4
IFNGR2	CRFB6	IFNGR2	1, 4
IFNLR1 (IL28RA)	CRFB14?	_	
IL10R2 (IL10RB)	CRFB4	_	5, 6
IL10R1 (IL10RA)	CRFB7	IL10R1	5, 6
IL20R1 (IL20RA)	CRFB8	IL20R1	7, 8
IL20R2 (IL20RB)	CRFB16	IL20R2	8
IL22RA2 (IL22BP)	CRFB9	IL22BP	3
TF	CRFB10 (duplication)	TFa	1
	CRFB11 (duplication)	TFb	1
IL22RA1	CRFB14?	_	
?	CRFB12	IFNUR1	This study
?	CRFB15	_	

Supplementary Table 2. IFNU and IFNUR1 genes in vertebrates

Species	Version	Gene	Coding Exons	Location or Accession Number	
Mammal					
Ornithorhynchus anatinus	Ornithorhynchus_anatinus-5.0.1	IFNUR1	7	NW_001794458.1: 89680908983283	
platypus)		IFNU	5	NW_001794186.1: 79796357986399	
Fachyglossus aculeatus	mTacAcu1.pri	IFNU	5	NC_052078.1:1199106211998896	
Australian echidna)		IFNUR1	7	XP_038604300.1	
Bird					
Inas platyrhynchos	BGI_duck_1.0	IFNUR1	7	NW_004677101.1: 470841475790	
mallard)		IFNU	5	NW_004676454.1: 459194462827	
Gallus gallus	GRCg6	IFNUR1	7	XP_015133867.1	
chicken)	N 1110	IFNU	5	CM000094.5: 1142565211429077	
<i>Jumida meleagris</i> helmeted guineafowl)	NumMel1.0	IFNU	5	NC_034410.1: 1079838010801894	
,	LICCO WEDTOI	IFNUR1	8	XP_021261921.1	
<i>agopus leucura</i> white-tailed ptarmigan)	USGS_WTPT01	IFNU	5	NW_024962334.1: 3965312239649735	
Cygnus olor mute swan)	bCygOlo1.pri.v2	IFNU	5	NC_049170.1: 1278068412784388	
Falco naumanni	bFalNau1.pat	IFNU	5	NC_054057.1: 4967790249673758	
lesser kestrel)		IFNUR1	7	XP_040470473.1	
Tyto alba	T.alba_DEE_v4.0	IFNU	5	NW_024881313.1: 2871875928714675	
Barn owl)		IFNUR1	7	XP_032851323.2	
thene cunicularia purrowing owl)	athCun1	IFNUR1	7	XP_026704474.1	
Coturnix japonica Japanese quail)	Coturnix japonica 2.1	IFNU	5	NC_029517.1: 1023957710242791	
<i>Torvus kubaryi</i> C.kubaryi_AGA036_p1.0 Mariana crow)		IFNU	5	NW_024581137.1: 4758454247580504	
Passer montanus ASM1480565v1 (Eurasian tree sparrow)		IFNU	5	NW_024356505.1: 3299600632991891	
Hirundo rustica Barn swallow)	bHirRus1.pri.v2	IFNU	5	NC_053450.1: 144857819144853799	
Serinus canaria Common canary)	cibio_Scana_2019	IFNU	5	NW_022041541.1: 1189395911898360	
Onychostruthus taczanowskii white-rumped snowfinch)	ASM1759005v1	IFNU	5	NW_024500349.1: 1454109214537005	
Pipra filicauda Wire-tailed manakin)	ASM394559v2	IFNU	5	NW_024104489.1: 135199139326	
Taeniopygia guttata zebra finch)	bTaeGut1.4.pri	IFNU	5	NC_044213.2: 1169336611697783	
onchura striata domestica Bengalese finch)	lonStrDom2	IFNU	5	NC_042567.1: 1194515411949189	
Catharus ustulatus Swainson's thrush)	bCatUst1.pri.v2	IFNU	5	NC_046221.1: 1153292711536898	
Centrocercus urophasianus Greater sage-grouse)	USGS_Curo_1.0	IFNU	5	NW_024884711.1: 3668861636685237	

Patagioenas fasciata monilis (band-tailed pigeon)	NIATT_ARIZONA	IFNU	5	LSYS01002888.1: 92230109226540
Limosa lapponica baueri (bar-tailed godwit)	Godwit_v1	IFNU	5	KZ505761.1: 1838121963
Meleagris gallopavo (turkey)	Turkey_5.1	IFNU	5	NC_015016.2:1649017516493601
Aquila chrysaetos chrysaetos	bAquChr1.4	IFNU	5	NC_044006.1: 6596426265960138
(golden eagle)		IFNUR1	7	XP_040975145.1
Dromaius novaehollandiae (emu)	droNov1	IFNU	5	NW_020452870.1:438128442620
Struthio camelus australis (African ostrich)	ASM69896v1	IFNU	5	NW_009271896.1: 40533714057941
Melopsittacus undulatus (budgerigar)	bMelUnd1.mat.Z	IFNU	5	NC_047527.1: 109446889109450503
Columba livia	Cliv 2.1	IFNU	5	AKCR02000004.1: 1170352611707123
(rock pigeon)	_	IFNUR1	7	XP_021145800.1
Apteryx mantelli mantelli (North Island brown kiwi)	AptMant0	IFNUR1	7	XP_013813805.1
Reptile				
Alligator mississippiensis	ASM28112v4	IFNUR1	7	XP_019337453.1
(American alligator)		IFNU	5	NW_017707901.1: 1517147915177353
Anolis carolinensis	AnoCar2.0	IFNUR1	7	XP_008119338.1
(green anole)		IFNU	5	
Chrysemys picta bellii	Chrysemys_picta_bellii-3.0.3	IFNUR1	7	XP_005301412.1
(painted turtle)		IFNU	5	NC_024222.1: 38797793885334
Alligator sinensis	ASM45574v1	IFNU	5	NW_005842092.1: 10219471027895
(Chinese alligator)		IFNUR1	7	XP_025061739.1
Crocodylus porosus (Australian saltwater crocodile)	CroPor_comp1	IFNU	5	NW_017728886.1 2044723420453945
,	GavGan comp1	IFNUR1 IFNU	7 5	XP_019391873.1
Gavialis gangeticus (Gharial)	GavGan_comp1	IFNU	3	NW_017729004.1:2664507126650909
Mauremys reevesii	ASM1616193v1	IFNU	5	NC_052624.1: 2497493924979882
(Reeves's turtle)		IFNUR1	7	XP_039345751.1
Trachemys scripta elegans (red-eared slider turtle)	CAS_Tse_1.0	IFNU	5	NC_048299.1:2409137424096381
Chelonia mydas	rCheMyd1.pri.v2	IFNU	5	CM026899.1:237610605237605534
(Green sea turtle)		IFNUR1	7	XP_027674429.2
Rafetus swinhoei (Swinhoes soft-shelled turtle)	ASM1942577v1	IFNU	5	CM033424.1:236101367236095006
Gopherus evgoodei (Goodes thornscrub tortoise)	rGopEvg1_v1.p	IFNU	5	NC_044323.1:2446451024469590
Chelydra serpentina (Common snapping turtle)	ASM1885937v1	IFNU	5	JAHGAV010000013.1:44785904473531
Pelodiscus sinensis Chinese_soft-shelled_turtle	PelSin_1.0	IFNU	5	NW_005851920.1: 952241945889
Varanus komodoensis (Komodo dragon)	ASM479886v1	IFNU	5	SJPD01000020.1: 1636026716369868
Podarcis muralis	PodMur_1.0	IFNU	5	NC_041323.1:1646223716468403

(Common wall lizard)				
Zootoca vivipara (common lizard)	UG_Zviv_1	IFNU	5	NC_048614.1:1490789214913259
Salvator merianae (Argentine black and white tegu)	HLtupMer6	IFNU	5	QVOM02000116.1: 4357996043569936
Sceloporus undulatus (fence lizard)	SceUnd_v1.1	IFNU	5	NC_056527.1:1585537415860670
Lacerta viridis (green lizard)	ASM90024590v1	IFNU	5	OFHU01004531.1: 280706288104
Aspidoscelis marmoratus (Marbled whiptail)	AspMar1.0	IFNU	5	MTQE01002638.1: 3617314636164211
Pogona vitticeps (central bearded dragon)	pvil.1	IFNU	5	NW_018150717.1: 9968492648
Sphenodon punctatus (tuatara)	ASM311381v1	IFNU	5	QEPC01013748.1: 89340318941185
Dermochelys coriacea (leatherback sea turtle)	rDerCor1.pri.v3	IFNUR1	7	XP_038271070.1
Amphibian				
Xenopus laevis	Xenopus_laevis_v2	IFNUR1	7	XP_018085133.1
(African clawed frog)		IFNU	5	MW924834
Rhinatrema bivittatum (two-lined caecilian)	aRhiBiv1.2	IFNU	5	LR584388.1: 112458824112468536
Hymenochirus boettgeri (Congo dwarf clawed frog)	UCB_Hboe_1.0	IFNU	5	CM033474.1: 3045208230458040
Xenopus tropicalis tropical_clawed_frog	UCB_Xtro_10.0	IFNU IFNUR1	5 7	MW924835 XP_002940450.1
Fish				
Latimeria chalumnae	LatCha1	IFNUR1	7	XP_005994806.1
(coelacanth)		IFNU	5	NW_005819652.1: 958231964496
Lepisosteus oculatus	LepOcu1	IFNUR1	7	XP_015193079.1
(spotted gar)		IFNU	5	NC_023187.1: 3931884839321474
Danio rerio	GRCz11	IFNUR1	7	NP_001035443.1
(zebrafish)		IFNU	5	MW547062
Callorhinchus milii	Callorhinchus_milii-6.1.3	IFNUR1	7	XP_007890021.1
(elephant shark)		IFNU1	5	NW_006890139.1: 28120742813536
		IFNU2	5	NW_006890139.1: 28229002825423
Astyanax mexicanus	Astyanax_mexicanus-2.0	IFNUR1	7	XP_007245285.2
(cavefish)		IFNU	5	NC_035906.1: 22081752210642
Polyodon spathula	ASM1765450v1	IFNU	5	NC_054536.1: 49552354957023
(Mississippi paddlefish)		IFNUR1	7	XP_041074472.1
Acipenser ruthenus (sterlet)	ASM1064508v1	IFNU IFNUD 1	5	NC_048325.1:53215715323376
	AmiCal1	IFNUR1 IFNU	7 5	XP_033870228.1
Amia calva (bowfin)	Amicali		3	CM030124.1: 4068671340688751
Anguilla anguilla	fAngAng1.pri	IFNU	5	NC_049204.1:3325208533253765
(European eel)		IFNUR1	7	XP_035288771.1
Anguilla rostrata (American eel)	ASM1855537v1	IFNU	5	CM031784.1:3338226733383949

Anguilla japonica	ASM635231v1	IFNU	5	VDMF01024909.1:38714033873144
(Japanese eel)				
Megalops atlanticus (tarpon)	MATL_1.0	IFNU	5	CM032876.1: 1225335312251769
Megalops cyprinoides (Indo-Pacific tarpon)	fMegCyp1.pri	IFNUR1	8	XP_036380572.1
Scleropages formosus golden arowana	ASM162426v1	IFNU	5	KV410874.1: 73617417363087
Asian bonytongue	fSclFor1.1	IFNUR1	7	XP_018610618.1
Heterotis niloticus (African bonytongue)	ASM1813684v1	IFNU	5	CM030905.1: 1117182611170081
Alosa sapidissima	fAloSap1.pri	IFNU	5	NC_055973.1:1703145017032575
(American shad)		IFNUR1	6	XP_041947255.1
Esox lucius	fEsoLuc1.pri	IFNU	5	NC_047589.1: 1581506315813807
(northern pike)		IFNUR1	7	XP_028977103.2
Clupea harengus (Atlantic herring)	Ch_v2.0.2	IFNU	4	NC_045168.1: 1822480618225835
Salmo salar (Atlantic salmon)	ICSASG_v2	IFNU	5	NC_027318.1: 1662506616626185
Oncorhynchus mykiss	USDA_OmykA_1.1	IFNU	5	NC_048575.1: 2519071325191829
rainbow trout		IFNUR1	7	NP_001117887.1
Chanos chanos (milkfish)	fChaCha1.1	IFNU	5	NC_044497.1: 1613468616135897
Pygocentrus nattereri (red-bellied piranha)	fPygNat1.pri	IFNU	5	NC_051213.1: 1053023110531859
Ictalurus punctatus	IpCoco_1.2	IFNU	5	CM004414.1: 3181609531812928
(channel catfish)		IFNUR1	7	XP_017315324.1
Tachysurus fulvidraco	ASM372403v1	IFNU	5	NW_020847794.1: 279125282142
(yellow catfish)		IFNUR1	7	XP_027034414.1
Cyprinus carpio	ASM1834038v1	IFNU	5	NC_056595.1: 45633764564202
(common carp)		IFNUR1	7	XP_018939179.1
Labeo rohita (rohu)	HRRL_Labeo_rohita_001	IFNU	5	JAFDUU010013128.1: 26892202688348
Colossoma macropomum (tambaqui)	Colossoma_macropomum	IFNU	5	NW_023494785.1: 89253538927409

Supplementary Table 3.

Primer sequences

Gene	Primer name	Sequence (5'-3')	Application
gapdh	qGAPDH-F	GTAACTCCGCAGAAAAGCCAGAC	qPCR
	qGAPDH-R	CAAAAGAAACTAACACACACA	
gig2	gig2-F	AAGATGACTTGGCTGGT	qPCR
	gig2-R	GATCTCGGCTGAGGTAGACG	
irf1	qDr-IRF1 F	GTGTCCAGAATGCGCATGCG	qPCR
	qDr-IRF1 R	GCCCACTGCTTGAACAGACA	
rsad2	zvig1.5	CGCCATCAGAGCATCCAGT	qPCR
	zvig1.3	TTCCACACCAACATCCAGAA	
mxa	qMX-F	AGACCATCCTCATTTCAGCAAACTCT	qPCR
	qMX-R	CAATCTTTTGTTGAATGAATCCCCTG	
ifnu	IFNUsgRNA-F	TGTAATACGACTCACTATAAAGACGTTCATT	sgRNA amplification
	C	GAGGCCCGTTTTAGAGCTAGAAATAGC	
	sgRNA-R	AAAAGCACCGACTCGGTGCC	
	IFNUmut-F	AGCAAAGAAAAGCAGTGTGG	Detection of ifnu mutant
	IFNUmut-R	CCAGATTTGGATTAAACCAAC	
	DRIFNMR4	CGCTGAAGAATCTCTCGCAC	RACE PCRs
	DRIFNMR5	GCCGTCTGACAGAGCGTTTC	
	DRIFNMF3	CAACTGTGCGTTTGACTATGA	
	DRIFNMF4	CCACGACACCTCCTCAATACG	
	DRIFNMF3-RT	CAACTGTGCGTTTGACTATGAT	qPCR
	DRIFNMR3-RT	CTCATTTTCAACACCGACCGAG	qr ore
	DRIFNM-EXF4	CGGAATTCCCACCATGGCCTGGATTAGAATCGT	Plasmid construction
	DRIFNM-EXR6	GGGGTACCTTAACGTCGCCGTCTGACAG	Tradifica Construction
ifnur1	DR12-EXF4	CCGCTCGAGCGACCATGGTGATGACTGGTTTTCTGACCTGC	Plasmid construction
illiui i	DR12-EXR5	CCCAAGCTTTTATACTTCATATTTGTTATTATG	i iusima construction
	CRFB12.5	CTGAATGGCTTCCAGGACAA	Knockdown test
	CRFB12.31	TGATGACTGGCGGAAACACTGAT	Kilockdowii test
	crfb12-E3-sgRNA1-F	TAATACGACTCACTATAGCCTACAGTCAGGACCTCTGG	sgRNA amplification
	choiz Es sgidvil i	GTTTTAGAGCTAGAAATAGC	sgreen unprincation
	crfb12-E3-F1	TCAGGGTAGCATTGTGACATCT	Detection of ifnur1 mutant
	crfb12-E3-R1	ATGGACCGTCTGACAAGTGAC	Detection of initial i mutant
crfb4	DRCRFB4-EXF1	CCGCTCGAGCGACCATGGTGATGTCCGCTTTTATCAGTTTTTC	Plasmid construction
C110 4	DRCRFB4-EXR1	CCCAAGCTTTTATATTATTTGCTGATTGTCC	i iasiniu construction
	DR4F3-MOtest	GAAGAATGTCCGCTTTTATCAG	Knockdown test
	DR4R3-MOtest	GCAACTTGTTTAGTCTCCACCC	Knockdown test
	Crfb4-B3	TAATACGACTCACTATAGGAGACTAAACAAGTTGCAG	sgRNA amplification
	C1104-D3	GTTTTAGAGCTAGAAATAGC	sgittiva ampimication
	Crfb4-F2	GGAGGGGTTATGATCCAGTT	Detection of crfb4 mutant
	Crfb4-F2	TAAGAGTATGCTACTGAAAACGTAT	Detection of crio4 mutant
crfb7	DR7-MOtest-F3	AACGAATCTTCCTGGTTGGTG	Knockdown test
CIID/		CTCACTGCAATAGATGTGATCG	Knockdown test
d . 0	DR7-MOtest-R3		V
crfb8	DR8-MOtest-F4	GGCAGGACACGGATCTCCCAG	Knockdown test
0.14	DR8-MOtest-R4	TGTTTGGGATGATCTTCCAGAG	IZ 1.1 · · ·
crfb14	DR14-MOtest-F5	TGAGTGCAGTGGTGACCGAAGT	Knockdown test
a 1 -	DR14-MOtest-R5	CTTCGCCACATACTCCACATC	77 11
crfb15	CRFB15.5	AGTGGACGTCTCCACACAA	Knockdown test
	CRFB15.30	CTTCCAGACTGTTATTCTGGAT	

crfb16	DR16F5-MOtest	GATCGGAGAAGAAGAGGTTATG	Knockdown test
	DR16R2-MOtest	GTGTTGAAGCCCACCTGAATAG	
xl-ifnur1	XL-C12RNAi-F1	TATGTGCGTGTCCCTTCCTTT	Knockdown test
	XL-C12RNAi-R1	TACAGTGCGATTCATTAGTCAG	
	XLCRFB12-F1	CTATGAAATGGCACAATGAGCAG	Gene cloning
	XLCRFB12-R1	GGATTGAGAAATATTGGGTCTG	
xl-il10r2	XLIL10RBLF2RT	AGCCTTCACTTGTCACCAGCAT	Knockdown test
	XLIL10RBLR2RT	ATGGCTTTGCTTAATTCTCCAG	
	XLIL10RB-F2L	AGCTCGGAGCACGTACCATTCC	Gene cloning
	XLIL10RB-R2L	GAGAAAAGCCCAGGCTGCCATTG	
	XLIL10RB-F1	GAGCACTTCTCATTCATTAAGCC	Gene cloning
	XLIL10RB-R1	GGGAAAAGCCGAGGCTGCAG	
xl-ifnu	XLIFNMLF1	AAATGCTGATGTCACAAATAC	Gene cloning
	XLIFNMLR1	TTTCAGGGTGAAGAAATGTTC	
	XLIFNMLEXF1	GGGGTACCCCACCATGCTGATGTCACAAATACTC	Plasmid construction
	XLIFNMLEXR2	GCTCTAGATCACTTGTCATCGTCATCCTTGTAGTCGGGTGAAG	
		AAATGTTCATGTT	

Supplementary Table 4. Morpholinos used in this study

Gene	GenBank ID	Sequence (5'-3')
crfb1	NM 001079681	CGCCAAGATCATACCTGTAAAGTAA
crfb2	NM_001077626	CTATGAATCCTCACCTAGGGTAAAC
crfb4	NM_001083868	GACATTGAAAAGTAACTTACTTCA
crfb5	NM_001034185	AGAGCGTATCCTCACCGTGTTTATC
crfb6	NM 001077627	AACTATCTCAAGTCCTTACCGATAT
crfb7	NM_001077625	ATGTCCACTGAAAAACCAACCTTAC
crfb8	NM_001079676	AAAATCCAAGGCAGTCACTCACTGT
crfb12	NM_001040353	ATACATATCTGAACTTACTGTCTCC
crfb13	GQ901864	TCTAAATGGTTTGTAGTTACCTGAG
crfb14	HM149255	CAGGAGAGATGATACTAACCTGTGA
crfb15	HM149256	AAAAACGGATTCACACTTACTGTGC
crfb16	HM149257	CTGAATCTGAAACGTGGGAACAATA
crfb17	GQ901865	TTAAACTAAATCGCCTTACCTTGTG
IFNphi1	NM_207640.1	CTGGTCCTCCACCTGTAATGCAATG
IFNGrel	NM_001020793.1	TTTCTGTGCTGTGAACCAAGTGATG
IFNG	NM 212864.1	TGAAGGCGTTCGCTAAAGTTAGAGT

Supplementary Table 5.

Genes used in sequence alignment and phylogenetic tree

Gene	Species, Gene Location or Accession Number
IFNAR1	human, NP 000620.2; chicken, NP 990190.1; zebrafish, NP 001029357.3;
	grass_carp, AGZ04472.1; tropical_clawed_frog, XP_002942842.5;
	platypus, XP_007667937.2; painted_turtle, XP_005283984.1; mouse, NP_034638.2
IFNAR2	chicken, NP_990189.1; mouse, NP_034639.2; human, NP_997468.1;
	zebrafish IFNAR2-1, NP 001073149.1; grass carp IFNAR2-1, AGW21650.1;
	zebrafish_IFNAR2-2, NP_001071094.2; grass_carp_IFNAR2-2, AMT92200.1;
	platypus, XP_028938459.1; tropical_clawed_frog, NP_001072391.1;
	painted_turtle, XP_042711707.1;
IFNGR1	chicken, NP_001123859.1; human, NP_000407.1; anole, AGU16999.1;
	zebrafish IFNGR1-2, NP 001165063.1; goldfish IFNGR1-2, ACV41809.1;
	zebrafish IFNGR1-1, NP 001129451.2; goldfish IFNGR1-1, ACV41808.1;
	platypus, XP_028913350.1; African_clawed_frog, XP_018120789.1; mouse, NP_034641.1
IFNGR2	human, NP 005525.2; chicken, NP 001008676.1; anole, AGL76447.1;
	tropical clawed frog, CX465469.2 and EL666073.1; mouse, NP 032364.1;
	zebrafish, NP 001071095.1; grass carp, AMT92201.1; platypus, XP 028937875.1
IFNLR1 (IL28RA)	mouse, NP 777276.3; chicken, K9JA28.1; tropical clawed frog, NP 001165238.1;
II (IEZOICI)	human, NP 734464.1; anole, AHY86485.1; whale shark, XP 020365568.1;
	elephant shark, XP 007892929.1; platypus, XP 028936126.1;
IL10R2 (IL10RB)	mouse, NP 032375.2; tropical clawed frog, NP 001165294.1; human, NP 000619.3;
ILIONZ (ILIOND)	chicken, NP 990188.1; anole, NP 001315154.1; zebrafish, NP 001077337.1 (crfb4);
	grass_carp, AHJ11133.1; common_carp, XP_018965682.1; catfish, XP_017326115.1;
II 10D1 (II 10D A)	platypus, XP_028937522.1
IL10R1 (IL10RA)	mouse, NP_032374.1; human, NP_001549.2; chicken, NP_001034686.1;
H 20D1 (H 20D A)	tropical clawed frog, XP_002932948.2; alligator, XP_014382148.1; coelacanth, XP_014345361.1;
IL20R1 (IL20RA)	human, NP_055247.3; mouse, NP_766374.1; chicken, XP_419723.1;
T 40D4 (T 40DD)	zebrafish, NP_001073144.2; trout, NP_001118088.1
IL20R2 (IL20RB)	human, NP_653318.2; chicken, XP_015150082.1; tropical clawed frog, XP_004917828.1;
	zebrafish, NP_001124447.1; catfish, XP_017351527.1; salmon, XP_013994619.1
IL22RA1	mouse, NP_839988.1; human, NP_067081.2; platypus, XP_001508953.2
	pigeon, EMC78530.1; zebrafish_CRFB14, NP_001184131.1;
	Sinocyclocheilus_IL22RA1L, XP_016374979.1; northern_pike_IL22RA1L, XP_010883684.1
IL10	human, NP_000563.1; chicken, NP_001004414.2; zebrafish, NP_001018621.2
IL22	human, NP_065386.1; chicken, NP_001186543.1
IL20	human, NP_061194.2; rat, NP_001137353.1
Type I IFN	human_IFNA2, NP_000596.2; human_IFNB, NP_002167.1; anole_IFN2, XP_016846269.1;
	anole_IFN1, XP_008101700.1; chicken_IFNA, BAA83089.1; chicken_IFNB, NP_001020007.1;
	tropical clawed frog_IFN1, CAO03085.1; tropical clawed frog_IFN2, CAO03086.1; carp_IFN3, ADI81045.1;
	carp_IFN1, ADI81047.1; zebrafish_IFNphi1, NP_997523.1; zebrafish_IFNphi2, NP_001104552.1;
	zebrafish_IFNphi3, NP_001104553.1; zebrafish_IFNphi4, NP_001155212.1;
	chicken IFNA3, XP 015132926.2; platypus IFN8, XP 028910681.1; platypus IFN7, XP 028910771.1;
	platypus IFN5, NP 001229647.1; platypus IFN4, NP 001229640.1; platypus IFN3, NP 001229639.1;
	platypus IFN1, NP 001229638.1; platypus IFN6, NP 001229650.1; platypus IFN2, NP 001229636.1;
	green anole IFN5 XP 016846269.1; green anole IFN3 XP 008101701.1;
	green anole IFN2 XP 008101702.1; green anole IFN1 XP 016846565.1;
	green anole IFN4 XP 008101700.1;
	African clawed frog IFNX1 to IFNX22, ANQ43313.1-ANQ43334.1
	African clawed frog IFN1 to IFN7, ANQ43306.1-ANQ43312.1
Type II IEN	
Type II IFN	human_IFNG, NP_000610.2; chicken_IFNG, NP_990480.1; zebrafish_IFNG, NP_998029.1;
	tropical clawed frog IFNG, ABU54059.1; anole IFNG, AGU16998.1; zebrafish IFNGrel, NP 001018629.1

	platypus_IFNG, NC_041741.1: 1683969616834260;
Type III IFN	human_IFNL1, NP_742152.1; human_IFNL2, NP_742150.1; chicken_IFNL, NP_001121968.1;
	tropical clawed frog_IFNL1, ACV32134.1; anole, AHY86484.1; platypus_IFNL3L, NC_041732.1:
	platypus_IFNL4L, NC_041732.1: 31948073191703;
	African_clawed_frog_IFNL1 to IFNL9, ANQ43335.1- ANQ43343.1;
	African clawed frog IFNLX1/IFNLX2, ANQ43344.1/ANQ43345.1

Supplementary Table 6.

Sequences of oligonucleotides for RNAi and probe

Gene	Application	Sequence (5'-3')
clawed frog IFNUR1	knockdown	CCTCATTCTTACTGTTACT
clawed frog IL10RB	knockdown	CATCTGAAAGAGTACCTAA
zebrafish IFNUR1	probe	TCGAAACATTGAGCAAGGCTGATGGAACGCG[tttCATCATCATCATCATCAT]30
	DIG labeled probe	DIG-ttATGATGATGATGATGT

Supplementary References

- 1. Aggad D, et al. In vivo analysis of ifn-gamma 1 and ifn-gamma 2 signaling in zebrafish. *J. Immunol.* 185, 6774-6782 (2010).
- 2. Aggad D, et al. The two groups of zebrafish virus-induced interferons signal via distinct receptors with specific and shared chains. *J. Immunol.* 183, 3924-3931 (2009).
- 3. Levraud JP, et al. Identification of the zebrafish IFN receptor: Implications for the origin of the vertebrate IFN system. *J. Immunol.* 178, 4385-4394 (2007).
- 4. Zou J, Secombes CJ. Teleost fish interferons and their role in immunity. *Dev. Comp. Immunol.* 35, 1376-1387 (2011).
- 5. Wei H, Wang X, Zhang A, Du L, Zhou H. Identification of grass carp IL-10 receptor subunits: functional evidence for IL-10 signaling in teleost immunity. *Dev. Comp. Immunol.* 45, 259-268 (2014).
- 6. Huo HJ, Chen SN, Li L, Nie P. Functional characterization of IL-10 and its receptor subunits in a perciform fish, the mandarin fish, *Siniperca chuatsi. Dev. Comp. Immunol.* 97, 64-75 (2019).
- 7. Lutfalla G, Crollius HR, Stange-thomann N, Jaillon O, Mogensen K, Monneron D. Comparative genomic analysis reveals independent expansion of a lineage-specific gene family in vertebrates: The class II cytokine receptors and their ligands in mammals and fish. *BMC Genomics* 4, 29 (2003).
- 8. Stein C, Caccamo M, Laird G, Leptin M. Conservation and divergence of gene families encoding components of innate immune response systems in zebrafish. *Genome Biol.* 8, R251 (2007).